

Pheasant / Gallinae / Grouse birds		Arunachal		NE India Sikkim		India		Elsewhere			
Species		Arunachal	Sikkim Bhutan	NE	India	E.L.	Altitude	Habitat			
Snow partridge Lerwa Lerwa		+	+	+	+	+			NW Pakistan, + S. Tibet, S. China		
Szechuan Sikkim snowcock T. t. aguilonifer		-	-	+	Sikkim				W. Bhutan		
Pheasant grouse											
Snow partridge Lerwa Lerwa		+	+	+	+	NW Pakistan S. Tibet, S. China	3000- 5000	Alpine meadows above treeline			
Sikkim snowcock Tetraogallus tibetanus aguilonifer		-	+	+	+	S Tibet	4500 5500	Alpine meadows rocky areas above			
Pheasant grouse Tetraophasis szechenyii		+	-	+	+	Upper subansiri + Siyom drainage of NEFA	3350 4600	rocky ravines with plenty cover in riv / rhodo forest			
Assam black partridge Francolinus francolinus melanostris		-	-	+	+	Assam Manipur	22000m	grassland Surma + Brahmaputra valleys			
Burmese francolin Francolinus pintadeanus phayeei		-	-	+	+	SE Manipur Kabaw + Manipur valley	?	low oak scrub in foothills. open up pterocarp forest			
Swamp partridge Francolinus gularis		-	Bhutan China	+	+	Assam W. / Bihar / Jy	-	Terrai grassland			
Tibetan partridge Perdix hodgsoniae		?	+	-	-	Tibet	3600- 5600	Tibet plateau			
Grey quail Coturnix coturnix coturnix		-	-	+	+	Manipur		Europe- Asia mixed			
Blue breasted quail Coturnix chinensis chinensis		-	-	+	+	Manipur, Assam		E - SE China Indo China Malaya			
Assam + Manipur painted bushquail Perdix perdix manipurensis inglisii / manipurensis		-	-	+	+	Assam NW Bengal NE Bhramaputra Assam Manipur		N / S India	< 1000m Elephant grass / swamps		
Common hill partridge Abrotophila torquata torquata		-	+	+	+	Extreme E. Assam N of Brahmaputra		N / S India Garhwal	SE Tibet 1500- 4000	ravines, khunds, hillsides of Banj oak etc.	
Assam hill partridge A. t. interstructa		-	-	+	+	Assam Sot Brahmaputra upper chindwin		Chittagon hills + hills Bavai, N. Cachar N Manipur	Upper Burma 2200- 2600	Wet temperate forest	
Kachin hill partridge A. t. batemani		-	-	+	+	? Hills of S Manipur		Burma Chin / Kachin hills	-	-	
Rufous throated hill partridge A. rufogularis rufogularis		-	+	+	+	Mishmi Hills		Kumaon- Nepal	-	1000- 2400	Lower than torquata Secondary scrub evergreen hill forest
Assam hill partridge A. r. intermedia		-	-	+	+	Sot Brahmaputra E Assam, Manipur H20 (Lushai hills)		Chittagon hills + hills Kachin	Upper Burma 600- 1800	Dense undergrowth & shady hills in evergreen forest	
White cheeked hill partridge Abrotophila atroregularis		-	-	+	+	Assam - (D. Brahmaputra Patla Mini, Mishmi) S. Sikkim, Naga Manipur, H20 tripura		Upper Burma E Pak	Plan - 7 1500 E Pak 7500	Wet undergrowth in evergreen forest & Bamboo	
Redbreasted hill partridge A. mandellii		+	+	+	+	Assam hills N of Brahmaputra		SE Tibet	350- 2450	Dense undergrowth in evergreen forest	
Assam bamboo hill partridge Bambusicola taylorii hopkinsonii		-	-	+	+	Manchhita - Patla Naga - Cachar, Khasi Jaintia Garo H20 Manipur - E Pak		Burma	< 2000	Open scrub in paddy - low oak & willow scrub	
Sikkim blood pheasant Ithaginis cruentus affinis			+			Sikkim W. Bhutan			2600 4600	Plenty in paddy & forest Birch, dwarf rhodo Vincet, juniper	
Tibetan blood pheasant I. e. tibetanus		+	-	+	+	Kanay + Subansiri		Western Nepal Black mtn of Bhutan	SE Tibet 4400	Dwarf rhodo scrub	
Mishmi Blood pheasant I. e. kuseeri		+	-	+	+	ANOR, Mishmi Luhit Neta		SE Tibet E -> Yunnan	2500- 4500	Rhodo scrub.	

**A rapid assessment of Biodiversity
in Mehao Sanctuary, Arunachal Pradesh
through Remote Sensing and
Geographic Information System**

Investigator

**S. Narendra Prasad
Salim Ali Centre for Ornithology & Natural History**

58.	Amachal	sikkim NE Bhutan	ME	EL.	Alt.	Habitat
Satyr tragopan Tragopan satyra	+	+	Darang in Assam Kameng Monjul?		2400 -4250	oak, deodar Rhododendron riyal Bamboo
Tibetan greybellied tragopan (3 specimens) Tragopan blythii molesworthi	+	E Bhutan →	NEFA Mishmi hills	SE tibet	1800 -3300	Thicket scrub & riyal undergrowth
Greybellied tragopan Tragopan blythii blythii	-	-	Assam S of BK Barail, Patkai Maza Manjor Mizo (Lushai)	Chin hills w of N Birma	71800	thick undergrowth in heavy evergreen forest
Temminck's tragopan Tragopan temminckii	+	E Bhutan (Subamir & Maha water w)	Luhit F.D. Mishmi hills upper Dibang & Tsampo	of NEFA S Tibet N Myanmar Szechuan	2100 -3500	dense undergrowth in evergreen forest on steep hillside.
Himalayan monal Lophophorn impejanus	?	sikkim E Bhutan	-	Burma?		
Sclater's monal L. sclateri	+	-	Kameng, Subamir Siang & Luhit	SE tibet NE Burma & Yunnan	3000 -4000	FR & Rhododendron forest.
Elwe's eared pheasant Crossoptilon crossoptilon harmani	+	-	Siang, Subamir & Luhit?	SE tibet Tsampo valley Salween valley	3000- 5000	grass slope edge on rhodo / sump scrub
Black backed khaleej Lophura leucomedana melanota	-	sikkim & w. Bhutan (Sankosh R.)	Dajediy	-	100- 2700	densely overgrown steep gullies thick undergrowth near streams & cultivation
Black breasted khaleej L. l. lathamii	+	E. Bhutan	Garo, Khai Cachar, Maza & Pathai	E - Prandy S - Muzh S - Arakan E - Dahuk	100- 2600	
Williams khaleej	-	-	? SE Manipur Mizo (Lushai)	Chin hills, lower & upper? Chinlain Arakan Yoma		-
Mottitt's khaleej	-	central Bhutan Isperlung	-	-	-	-
Red jungle fowl gallus gallus unghii	-	(West Bengal Assam)	→	-	-	moist deciduous forest
Burmese red jungle fowl G. G. spadiceus	+	-	Eastern Mishmi hills NEFA	sw Yunnan Burma, Laos Malaya, Hainan	-	
Mrs. Hume Barred backed pl. Sympterus himiae himiae						

A rapid assessment of biodiversity in Mehao Sanc
Arunachal Pradesh through remote sensing and Ge
Information System

Investigator

S. Narendra Prasad

Salim Ali Centre for Ornithology & Natural History

Kalampalayam, Coimbatore-641 010