

- Falco Jaffer-Laffer* - not abundant
Lit-hofaleo chicquena - ~~Tam~~ Turumbi V. Rare
 Nestil - abundant
 Shivra - local distribution
 sparrow Hawk - Borabhum, March, Tichele
Spizaetus Lalthami - Hawk Eagle - killed at Sisda Borabhum (for
 detail see II) 1 seen in Lucia
 Rufous-bellied H. Eagle - Chaibansa - T. (detail Vol I - 316)
Circusetus gallicus - Serpent Eagle - very probably occurs.
Spirorthis chela c. " " Common
Circusetus leucogaster - Grey-bellied Sea Eagle - no means rare - found on large rivers
 Pale Harrier - in all the open parts
 Montague's Harrier - not common
 Pied " - not very common - parts of division
 Marsh " - common in the cultivated parts
 Marsh with - found
 Common " - abundant
Milvus major - Large with - occasionally observed. Lohandagga. sp.
 erect-ed Honey Buzzard - sparingly - sp. Lohandagga
 Rajmahal Hills appeared to be common
 Black-winged with - not common in c-pure.
Strid indra Screech owl - occasionally
Epiaur candida - Frequents the long grass jungle. T.
Asealaptra -
 Rock-horned owl - Western c-pure
 Dusky-horned " - not common
 Brown Fish owl - occurs

Scops pennata - Scops owl - 1 believed, only sparingly in Epure
Spotted owl -

Heteroglaux Brewitti - should be looked for in Epure (Mr. Hume)

Athene radiata - Jungle owl - not uncommon

Ninox hirsuta - Indian Hawk owl - not common - sp. near Kaveri - one sp. by T. in Dalbhum.

Common Swallow - Common.

Wire tailed " - not frequently met

Red-rumped " - sometimes seen in vast flocks.

White-rumped spine tail - near Chenbasa (T.)

Cypselus melba - Arabian swift - Singhbhum (T.)

Common " - " Rare in Maunbhum, although seen in Puna bia itself (Capt. Beavan)

Palm Swift - In abundance where its favorite trees are common.

1. Crested Swift - parts of Epure - but nowhere abundant Rajmahal hills also.

Caprimulgus indicus - night-jen - Lohardagga -

" albonotatus - Large bengal N.J. - T & Beavan obtained it. Ball

" asiaticus - 1. N. Jen - only heard & seen - abundant in Maunbhum (Beavan)

Franklin's N. Jen - Rajmahal (Ball) - abundantly distributed - also Rajmahal

Harporctes fasciatus - Malabar Trogon - Dampara in Dalbhum T. single sp.

Ball says: this bird does not appear to have been obtained or even seen by any subsequent observer in Epure.

C. Indian bee-eater - abundant. Beavan says - it breeds in Maunbhum at the begin ning of April.

Black-tailed bee-eater - very rare. only met it in the Western part of the Division March April.

Chestnut headed bee-eater - one pair in Epure

1. Rooster - Very common - breeds in the station
of Ben Coorah in April (Beavan)

Brown headed kingfisher - Also Rajmahal

White breasted N.F.

Common kingfisher - v. c.

Ceryx hidacetyla, Pallas, has been seen neighbourhood of the Noel river in Palamanu

Alcedo asiatica does not occur in the Dinivim. in the Brahmani river in the
Rajmahal - shot in 1869.

Pied kingfisher - open parts

H. *coronata* - Malabar Pied Hornbill - affects estate beaches in Epure
6 to 10. shot in Manbhum, S'bhum shot
in the wood from Nanchu' plateau to Purukhi

H. *albinostis* - (See p. 388 II)

Common Hornbill - Manbhum, S'bhum, A'burgh. Rajmahal

Alexandrine Parakeet - heavy gurgle. Rajmahal. Common.

Rose nighed P. Common. (a nest with 3 eggs at

Pranurria in Manbhum on the 1st March 1865. Beavan)

Rose headed P. Common

Picus maharattensis - (See p. 390 III)

Southern Pig my Wood pecker - Sirsuga.

Chrysocolaptes Sultaneus - Golden backed Wood - (391 II)

Black backed Wood pecker - Palamanu. shot-3

Geopelia striolata - Small green Wood p. Rare in Epure. Collections confined
to the Western part. Lohardagga

Micropternus thaliocephalus Bengal. Rufous Wood p. - Manbhum Beavan (Rajmahal Bull.)

Brachypternus aurantius - Y. Backed - Occasionally in the forests of Epure

Yund forquilha - C. Wrench - obtained in Manbhum (Beavan) Rare
bird in Epure. (Bull, Palamanu. Rare (Levin)

C. Green barbet - Also Rajmahal hills.

e. B. Parrot - Common

Cuculus canorus - European cuckoo - Lohandagga - Ranchi station. M. bhum

c. Hawk cuckoo - throughout e. pure. Chot - M. bhum, S. bhum, B. bhum and Rajmahal.

1. Palearctic cuckoo - occurs sparingly. Manbhum (Beavan)

Pied bellied cuckoo - Very rare in e. pure. Saw one in eastern part of M. bhum and chot in B. bhum Palamau (Levin)
(More frequent by
Noel. Common. Seldom met in more jungle part

Zamelostomus bistris - Large green billed Malkoha. (See p. 394)

Crow pheasant - (See p. 394) Rajmahal hills

Centrocoryx bengalensis - Indian Coucal - (See p. 395)

Trochona sirhee - Bengal sirkeer - Palamau - Rajmahal

" *affinis* - Central I. " - (See p. 395) + 396) sp. Palamau (Levin)

Aethopyga miles - Red honey-sucker - B. bhum + Between Ranchi +

Purukh. exact on the spot east of Jona. (Hall, 396)

Leptocoma zeylonica - Amethyst-rumped honey-s. S. bhum T. Manbhum Beavan.

Breeds in March - April

Purple Honey s. Common in e. pure.

+ Flower pecker - Common

thick-billed flowerp. - equally distributed in e. pure.

Salpornis spilnotata - spotted grey creeper - in 1868 near the Thakda hills in M. bhum. Kaudera in Lohandagga.

Chesnut-billed nuthatch - Rather rare in e. pure.

velvet-fronted nuthatch

Upupa epops - European Hoopoe - pretty generally distributed throughout the open parts of e. pure - Manbhum (Beavan)

Grey & Lactora - Grey shrike - Rare in Manbhum (Beavan) Hall did not see this species. Palamau (Capt. Grey A. C. Lohanda 1912)

Collyrio tephronotus - Grey backed shrike - Not common. Rajmahal abs.

C. nigriceps - Black headed shrike - tolerably common. abundant Rajmahal

- Lanius vittatus* - Bay-backed shrike - occasionally seen Mbhum (Beavan).
Same is the case in the eastern parts of the dist., but
common at Palamanu + H. Bagh (B), Rajmahal.
- L. cristatus* - Brown shrike - obtained in Mbhum + Sbhun. Common everywhere.
- C. Wood shrike - Common (^{obtain} ~~found~~ see p 399)
- Hemipus picatus* - Little Pied shrike - Very rare. only Sbhun
- Votroivora melaschistos* - Dark-grey cuckoo shrike - Rare. Manbhum (Beavan), Sbhun +
Mbhum + Lohandagga (Ball,
- V. sykesii* - Possibly occurs in the eastern parts of the division. Rajmahal
↓ shot in Palamanu (Levin)
- V. melaschistos* - Rajmahal
- Granaealus macei* - Large cuckoo shrike - H. Bagh, Mbhum + Lohandagga.
(See p. 400)
- Large Mini vet - Throughout Empire. Common in Mbhum (Beavan)
- Small " - Common. abundant in Lohandagga.
- Orange - Common.
- x *Buecharya longicaudata* - Grey drongo - thin seen in Mbhum. Obtained Rajmahal
White-bellied drongo - throughout Empire.
- Large Racket-tailed drongo - Very rare. Common in Mbhum (Beavan)
Rajmahal hills. Common in the extensive bamboo jungle of Divi (Beavan)
- Hair-curtled drongo - Manbhum (Beavan), Main Pat in Sirsuga (Ball,
T. collected within our limits. (403)
- Army swallow shrike - Rare. Rajmahal hills also (291 III)
- (292 III) { P. flycatcher - obtained at Nashurpurk in Mbhum (Beavan),
Lohandagga B. Beavan is of the opinion that it only come to Mbhum
"When the new leaves come in to the trees at the end of March".
Baleh-naped blue flycatcher - obtained no part of the
division. "In Mbhum it is seen frequently in suitable localities
(Beavan) in Jan and Feb, but does not apparently make a lengthened stay".

- Zenaidura macroura* White-throated Fantail fly - Not often seen in Epure
 White-browed " " Not-uncommon in the mango grove
 Grey-headed flycatcher - Occasionally found in mango grove.
 but they prefer the neighborhood of rivers with well-wooded banks.
- Alcedo latirostris* - Included in Blyth's list of T's birds. Bull never obtained it.
 Verditer flycatcher - Not very common in any part of Epure
 Bull generally met it near rivers in little frequented jungles.
 Tichell's blue red-breasted fly - Rare. sbhum
 White-browed blue fly - Rarity - sbhum T, dohardagga (Ball, Dgunj
) (Sep. 292 III) Levin
- Muscicapula ruficeps* - Obtained in Mohum Beavan, ?
- Erythrostera albicilla* - White-tailed Robin fly - Common
 Red-breasted fly - shot several sbhum in Jan 1869
- E. Puzilla - Rufous backed fly - a specimen. sbhum Feb 1869
 little pied fly - sbhum T, Mohum Beavan (B)
 Malbar Whistling thrush (page 406 also 292 III)
 " " 407 also 292 III
- Pitta coronata* - Yellow breasted ground " - " 407 also 292 III
 Blue Roen thrush - Near mouth of the large rivers. shot at sbhum,
 Hbagh.
- White winged ground th - Rare in Epure.
 Orange headed gr. th. - Mohum Beavan, sbhum T, Rajmahal B
 Darky ground th. sbhum T, Rajmahal Bull (P. 408)
 Small-billed Mountain th. - Mohum Beavan, occasionally Western
 Epure. Rajmahal as late as April.
- Yellow-eyed Babbler - Throughout. Common in Palamanu.
 Yellow-breasted babbler - Borabhum + Dhalbhum T,
 Banks of the Cosye (Beavan,
 Rufous billed Babbler - Kerurgurh in Mohum Beavan,
 Spotted babbler - Rajmahal. (T & Beavan also) (409,
 Southern Scimitar babbler - should be looked for as it
 probably occurs.
- Bengal babbler - Common. Breeds in Mohum Beavan,
Actinops caudata - striated babbler - Rather common in Western Hbagh & Palamanu
 Beavan did not obtain in Mohum.

White throated bulbul (p. 410)

Black eyed yellow bulbul - obtained by T, but by no one since in epure. He says "I frequented the beautiful hanging woods of Dampana in Dholbhum, when also alone I met with him."

Red whiskered bulbul - Eastern parts of Mbum, but rare. Not seen it in any part of the parts of the Division.

Pyrenonotus chrysorrhoeus - Madras bulbul (p. 410) also 292 III

P. Terdoni - Green bulbul - Abundant. Barahen, Sbum, Hbagh + Rajmahal
- Gold-fronted G. bulbul - less common. (411)

Aegithina tephra - (411) not uncommon in epure. Sbum, Hbagh + Rajmahal
Indian Oriole - Probably sparingly throughout Divi.

Mbum (Beavan), Rajmahal. Unidentified.

Golden Oriole - Abundant, Rajmahal III

Magpie Robin - Occurs

See 293 III (Shama - extremely rare. once in 2 days.
Mbum - (breeds in April-May Beavan), Rajmahal
Indian Robin - very common. (p. 412)

Pratincola caprata - White winged beach Robin - cannot be very common in epure.
Borobhum T, 413,

" *indica* - 1. bush chat - tolerably abundant.

P *leucura* - Myth, may v. possibly occur in epure.

Brown Rock chat - T, 4 ymores at Rotasgarh. Ball.

Red start - common. Rajmahal.

Ruby throat - Dampana in Dbum T. (413)

Blue throat - Rare. Mbum B Palaman (Loring)

Lesser Reed Warbler - (114)

Tailor bird - Not common.

Franklin's Wren Warbler - common in most parts of epure

Mbum Beavan,

Long Tailed W. Warbler - Barabhum + Dbum T,

Dymipus neglectus - T, no doubt it occurs.

- D. inornatus* → E.W. Warbler & Palamanu (Lavin),
 Rajmahal also epure → Indian Museum Catalog
 Skyes Warbler - M bhum & S bhum by the people.
- Phylloscopus tristis* - Brown tree warbler - occurs but not common. M bhum,
 S bhum and Lohradaga.
1. *Fuscatus*, *megubris*, *nitidus* & *viridanus* (Beavan believes occurs)
4. *affinis* - Tickell's tree warbler - S bhum, & other parts of Dini
 in 901-
- Reguloides proregulus* - Crowned Tree W - Palamanu also Satpuras.
- Abroornis cantator* - Linn's black-browed W - T, but has escaped subsequent
 collectors in the Dist - when it was first obtained
- Sylvia Terdoni* - Large black-capped W. Nakhur surh in M bhum (Beavan),
 Bull did not see it in epure but frequently met with in the
 Nambadda valley at the foot of the Satpura hills.
- M. madraspatana* - Pied Wagtail - Common. (Cosey river Beavan)
- M. luzoniensis* - White faced Wagtail - Common. Specimen H bugh, S bhum,
 Mambhum.
- M. dukhunni* - one specimen of what I take to be the Black faced from
 epure & Rajmahal.
- Calobates melanops* - Grey & yellow Wagtail - uncommon in streams passing
 through jungle.
- Budytes flava* - Common
- B. citreola* - Yellow headed also occurs. (S bhum Bull. Common in the Lohji & Norel rivers
 in Palamanu during local W. 1873-74 Lavin)
- A. arboreus* - Tree pipit - Common
- A. meenlatus* - S bhum + Barahan.
- Corydalla Richardi* - Large marsh pipit - At Assensoh (416)
 Indian Museum ^{Cat} S bhum.
- C. rufula* - 1. tit-lark - very common.
- C. striolata* - Large tit-lark - abundant. Rajmahal
 White eye - somewhat sparingly occurs. M bhum Beavan
 Grey tit - occurs sparingly throughout Chentans T.
- Macrotrochus Terdoni* - Southern yellow tit - occurs sparingly throughout. Also
 six Saja: Satpura hills common Jan + Feb.
- M. Karthogeny* - does not occur in these jungles.

Jungle crow - spread throughout C. pure

Common - Crow is very common

V. rufa

Tree pie - throughout division. Beavan says rare in M. hum. (detail p. 418)

Pied Myna - tolerably common except heavy jungle

C. Myna - abundant

Bank n - rare. actually obtained only in S. hum

Aeridotheres fuscus - Cattle Myna - does not occur or if does only sparingly in the eastern parts of the division. In Sirgaja common.

Black headed Myna - sparingly spread throughout the Division. Abundant in Sirgaja.

Grey headed Myna - less common. obtained only in M. hum and S. hum, Bal,

Race-colored Starling - may be looked for with certainty in Feb. wherever there are cotton trees. Large flocks roosting on the islands in the lake at Purneha as late as April (Beavan)

S. hill Myna - Jungles of Saranda (p. 420) also 293 VII

Common Weaver bird - throughout C. pure.

P. manyan

- may very possibly occur.

Spotted Munia - sparingly - Rajmahal also.

(White-backed Munia - S. hum. Marbhum (Beavan), ~~White~~ 420)

Plain brown Munia - obtained in M. hum. observed in Sirgaja and other parts, but I do not think it is common anywhere in the Division.

Red Munia - All parts of the division.

Green Munia - certainly occurs in Sirgaja, if not in other parts of the division. No means rare in the Satpura hills and the Nerbada valley.

H. sparrow - common.

Cymnoris flavicollis - yellow-necked sparrow - M. hum + S. hum and Sirgaja. But no where so abundant as it is in the Satpura hills, when I hardly passed a day without seeing numbers.

e. Rose Finch - is not abundant in epure. Obtained in both sbhum + Sirsujā. With former dist- abate at the end of April.

Bergal bush lark - not uncommon in Mbhum (Beawan), one of this species by Ball.

Mixis affinis - this is the common species (421) abundant in sbhum Red winged bush lark - occurs in Mbhum (Beawan) in Palaman. Ball.

Singing bush lark - sbhum (T)

Rufous Tailed Finch l. abundant in sbhum + Palaman

Black-bellied F l. Very common

Short-toed lark - common in Mbhum (Beawan)

specimens killed at Kaskur gush have a large blackish patch in each side of the breast - above. At-

sirsujā - vast flocks in March - April on the grassy plains.

Alcedo Gularula - very possibly occurs, but neither Beawan nor Mysore obtained it.

Green pigeon (for local obtained p 423)

orange-breasted green pigeon - "in some abundance near Ambekhanuggur (Mbhum) in Dec. 1864" (Beawan)

Imperial pigeon - very locally in epure (for detail 424)

Purph Wood pigeon - sbhum (T) Mbhum, on the banks of the Coryst near Ambekhanuggur (Beawan, Sirsujā Ball)

Blue rock pigeon - very common. (425)

Rufous Turb Dove - not common in all parts. Mbhum Beawan most common in Sirsujā Ball, also Rajmahal.

Little brown dove - occurs throughout. Mbhum (Beawan)

Spotted dove - throughout but not so common as the preceding.

Ring dove - throughout.

Red Turb dove - throughout. Abundant in Sirsujā.

Bronze-winged Dove - extreme rarity in epure. Seen once in the forest in the tributary stāk of Gangpur Khat

II

Rajmahal. obtained by T in S. Ghum.

Painted Sand grouse - Western parts of the division,
Surguja & Palamanu. Breeds in Palamanu according to Levin.
Common Sand grouse - Palamanu.

Peacock - abundant in all the hilly parts of Epure.
(See p. 426)

Red Trough fowl - abundant (See p. 426)

Hepburnia spadiceus - (p. 426) Red sp. fowl's skin which was shot at Epure. Bone removed (Bell)

Sp. fowl - more common. M. Ghum (Beavan),

Black Partridge - Common. Abundant in the West - when few 'Grey'
are to be met with - M. Ghum, Beavan, rarely, Palamanu shot Levin

Grey Partridge - most parts but rare in the extreme West -
abundant in Palamanu

B. sulavis - Nymph - does not occur in Epure. Abundant in low jungle
bordering sheets to the east & north-east of the Rajmahal hills.

Trough back Quail - M. Ghum, Beavan, possibly most common

Rock back " - occurs. one female sp. from Palamanu (427)

Microperdix Mewilli
p. 294 III

Painted b. Quail - shot in Western Epure - Surguja also Satpura

Blue breasted Quail - occurs in Epure.

(Turnix taiqoor - Mach-breasted banded Q. " " " " not in common M. Ghum (Bea
294 - III
Button Quail - tolerably common - Abundant in the vicinity of

Panastath (Beavan), (428)

Synaeoides aurota - Linn & Florinheim - (See p. 428-29)

Cuvierius Coromandelicus - Courcier's Plover - (429)

Eporeola lactea - Small Swallow Plover - obtained Ramdola river in M. Ghum
by Beavan & Bell. Abundant in the zone, when the river
boards Epure in the north.

Golden Plover - rare occurrence in Epure.

Aegialitis Philippinus? Common in the larger rivers. Breeding in M. Ghum in
March (Beavan) Also Rajmahal hills. (429)

Red-Wattled Lapwing - most common birds in Epure

Yellow " " - Rare, met with in all parts of the
division.

12

(Spar - winged Lapwing - I have never seen in Epure.

Oedieremus indicus - Beavan says: *S. bilobus* is rare, and this bird not so common in Mblum. According to him this is by far the rare bird of the two.

He seen it in Mblum, Sblum & Sirgija. (430)

Sarus crane - rare Mblum (Beavan) No. 1. Common in the open valleys of Sirgija (430)

Common crane - Large flocks high up in the air. Clearing of Bains is no uncommon sound in the parts of the division.

Frequently seen in the vicinity of the Damodan & Mahan rivers.

Demoiselle crane. Not seen in Epure.

Wood snipe - Not much to be seen in the division. Flashed once in the beginning of March at the foot of some hills in the Spara of Sirgija.

Gallinago stenura - Pin-tail snipe - "A fine specimen killed by me at Ambekam nggan in Mblum was larger than stated by Dr. Soden ----"; Beavan, not in Epure Ball.

C. snipe - occurs in suitable localities.

Teeu snipe - no personal knowledge of the occurrence in Epure.

Painted snipe - found throughout (431)

Little stint - only species of *Fringa* which he collected in Epure.

Spotted Sandpiper - in all the rivers of Epure

Green " " - Same.

Common " " - Same.

Green shanks - Common in Epure.

Bronze-winged Terebra - Not common

Pheasant - Lebed " - occasionally in fields in Epure

Purple Coot - Same.

Bald Coot - Rather common in tanks & fields.

Gallinula chloropus - Water hen - When there are fields & tanks.

White-breasted Wren - Occasionally met with.

Adjuvant. (Forbill tail 432)

Cranopolarhus javanicus - Hair-crested crane - Mambum + Sirgija

- Nyctenia australis* - ¹³ Rare in Epure. (433)
 Black stork - (433)
 C. White stork - Several parts of the division. (433) &
 six gaja.
 White-necked stork - abundant. (433)
- Ardea cinerea* - Blue Heron - not common but found throughout
 Purple Heron - Throughout
- Herodias alba* - Not very common
 " *intermedia* - Common enough in the well-watered parts. Rare in the West.
 " *garzella* - See remarks on last species
 Cattle egret - in all parts. (434)
- Ardeola leucoptera* - Common in all parts
 Little green heron - in all most every river.
- Ardeola cinnamomea* - (435)
 Night heron - Common. (435)
- Scolopax leuccephalus* - Rare in Epure. Seldom seen it. Killed one in the Noel river
 Patnamau.
- Anastomus oscitans* - shall ibis pretty generally distributed. only on one
 occasion by Beavan. Rajmahal
 White ibis - All parts of the division. (436)
 Black ibis - Common. Generally in flocks.
 Glossy ibis - Very rare. only once saw a small flock
 of them in the wing. ~~in~~ In the neighbourhood of the Ganges close
 to Rajmahal is common.
 Flamings - it may very possibly occasionally occur there.
 Grey goose - Rare visitor.
 Banded-headed goose - once shot in Epure in the
 Damodar river, Dec. 1872.
- Sarkidiornis melanotos* - Black-backed goose - very common in the Western parts (437)
 Cotton tail - in all parts when there are tanks.
 Whistling teal - in all parts. (437)
 Brambling duck - common in large rivers. Damodar, Goppe,
 Sukam nihha, Res, Mahan etc. (437)
 Shoveller - Rare. At Rajmahal
 Spotted billed duck - "rare, in pairs" (Beavan, extreme
 rarity in all parts) (Rare pair shot in the head of the Goppe

14
rivers in Sirgupa.

Pink-headed duck - never seen in any part of the division. Others who met it in M. Bhum.

Gadwall - found whenever there are plenty of tanks & sheets.

Pink-tail duck - found with the Gadwall in the larger rivers. Large flocks in M. Bhum but Beasem did not see this bird.

Common teal - found in all parts of the division but not in great numbers.

Blue-winged teal - less common. one year B observed that it did not appear in M. Bhum until after Jan.

Red-necked Pochard - not rare in M. Bhum & some of its parts where tanks occur.

Aythya ferina - Red-headed Pochard - shot in M. Bhum by Mr. Wilcox, S.P. in M. Bhum. But Ball never met this bird.

White-eyed duck - frequently in the well-watered parts.

Tufted Duck - in large flocks, not uncommonly.

Mergus castor - (Sep. 439)
ditch grebe - in all parts.

Hydrochelidon indica - (439)

Black-billed Tern - in all the principal rivers of Upper

S. Beena, Sykes.

seen in some of the larger rivers, but have no record of it that we can find.

Rhynchos albicollis - shot in the Ganges near Rajmahal.

Large Cormorant - occasionally in the larger sheets, as in the Govin Shatta in Sirgupa.

Less Cormorant - not very common.

ditch Cormorant - very common.

crane bird - singly or in pairs in rivers throughout division.

- Euphonia Bonellii* - Curtius Hawk - Engl - Palamau - Mr. Levin of Dattargung
- Uromitis filifera* - (nest - in April in a cleft - in a rock in the Noel (detail 289-III)
- Acanthylis sylvatica* - by Levin. Abundant on the southern frontiers of epure Ball
- Cypselus affinis* - e. Indian Swift -
- Merops philippensis* - small parties on the banks of the Noel in Palamau
(Levin)
- Alcedo asiatica* - (p. 289 vol IV)
- Hydrocissa albirostris* - shot at Lobloi near Lanigarh in the south of the division.
small party in an old grove of mango trees. In the same
Noel of County he also met with *Coronata*, two species
overlap each other's limits (Ball,
- ~~*Myra torquilla* - Wryneck - obtained in Palamau (Levin) very rare.~~
- Pericrocotus erythropygus* - White-bellied Minoret - (See p. 291 vol III)
- Painted Partridge - shot by Levin in Palamau.
- Esacus recurvirostris* - Large-stone Plover - Noel river in Palamau Levin. Parties
in the very beds of the Mahanadi + Ebe in the Sambal
pur dist - Ball.
- Hydrochelidon indica* - small Marsh Tern - Common in the Noel in Palamau
(Levin)
- Sterna Seena* - Large river Tern - Noel - Levin
- Pelecanus javanicus* - Pelican - obtained + says is far from rare. Levin